


FORM PTO-1449

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICEATTY. DOCKET NO.
NIH171.001C1APPLICATION NO.
10/005,305INFORMATION DISCLOSURE STATEMENT
BY APPLICANT

(USE SEVERAL SHEETS IF NECESSARY)

APPLICANT
Wang, et al.FILING DATE
November 2, 2001GROUP
1646

U.S. PATENT DOCUMENTS

EXAMINER INITIAL	DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE (IF APPROPRIATE)

RECEIVED
 JUN 06 2002
 TECH CENTER 1600/2900

FOREIGN PATENT DOCUMENTS

EXAMINER INITIAL	DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION	
						YES	NO
Book	1. WO 96/40191	12/19/96	PCT				

OTHER DOCUMENTS (INCLUDING AUTHOR, TITLE, DATE, PERTINENT PAGES, ETC.)

EXAMINER INITIAL	OTHER DOCUMENTS (INCLUDING AUTHOR, TITLE, DATE, PERTINENT PAGES, ETC.)
Book	2. Brown, L. E., et al. (1995) Synthetic Peptides Representing Sequences Within gp41 of HIV as immunogens for murine T- and B-cell responses. Arch. Virol. 140(4):635-654.
	3. Kilby, J. M., et al. (1998) Potent Suppression of HIV-1 Replication in Humans by T-20, a Peptide Inhibitor of gp41-mediated Virus Entry. Nature Medicine 4:1302-1307.
	4. Lawless, M. K., et al. (1996) HIV-1 Membrane Fusion Mechanism: Structural Studies of the Interactions between Biologically-Active Peptides from gp41. Biochemistry 35(42):13697-13708.
	5. Su, S. B., et al. (1999) T20/DP178, an Ectodomain Peptide of Human Immunodeficiency Virus Type 1 gp41, Is an Activator of Human Phagocyte N-Formyl Peptide Receptor. Blood 93(11):3885-3892.
	6. Su, S. B., et al. (1999) T20/DP178, an Ectodomain Peptide of Human Immunodeficiency Virus Type 1 gp41, Is a Potent Activator of Human Phagocyte N-Formyl Peptide Receptor. FASEB Journal 13(4):pA293 (Annual Meeting of the Professional Research Scientists for Experimental Biology - April 17-21, 1999).
	7. Su, S. B., et al. (1999) T21/DP107, A Synthetic Leucine Zipper-Like Domain of the HIV-1 Envelope gp41, Attracts and Activates Human Phagocytes by Using G-Protein-Coupled Formyl Peptide Receptors. J. Immunology 162(10):5924-30.
	8. Tas, M., et al. (1988) A Monocyte Chemotaxis Inhibiting Factor in Serum of HIV Infected Men Shares Epitopes with the HIV Transmembrane Protein gp41. Clin. Exp. Immunol. 71(1):13-18 (Database Biosis Online, Biosciences Information Service, Philadelphia, PA 1988).
	9. Wang, J. M., et al. (1999) T20/DP178, An Ectodomain Peptide of HIV-1 gp41, Is a Potent Activator of Human Phagocyte N-formyl Peptide Receptor. AIDS Pathogenesis, Keystone, CO, January 7-13, 1999.

O:\DOCS\MXG\MXG-1349.DOC:vb.052802

EXAMINER <i>E. Kemmerer</i>	DATE CONSIDERED <i>6/30/03</i>
*EXAMINER: INITIAL IF CITATION CONSIDERED; WHETHER OR NOT CITATION IS IN CONFORMANCE WITH MPEP 609; DRAW LINE THROUGH CITATION IF NOT IN CONFORMANCE AND NOT CONSIDERED, INCLUDE COPY OF THIS FORM WITH NEXT COMMUNICATION TO APPLICANT.	